



To whom it may concern

Dr. Stefan Dreiheller
Quality Unit

Date of issue: 23.Januar.2023
Page 1/3
Doc.-No. EIP-3036
Revision 7

Nitrosamines

EMA/369136/2020; FDA Guidance for Industry „Control of Nitrosamine Impurities in Human Drugs“

MEGGLE Product: MicroceLac® 100

The formation of potent genotoxic nitrosamines as impurities is possible in the presence of secondary, tertiary, or quaternary amines and nitrite salts under acidic reaction conditions.

Regarding the requests of EMA-CHMP and FDA-CDER that marketing authorisation holders for human medicines containing chemically synthesised active substances have to evaluate the risk for the possible presence of nitrosamines (e.g. NDMA, NDEA, NMBA, NMPA, NIPEA, NDIPA), a risk evaluation was conducted for the MEGGLE product to support the marketing authorisation holders. This evaluation is based on the [IPEC Questionnaire](#) and attached.

The product is a co-processed, directly compressible spray agglomerate comprising 75 % Lactose Monohydrate (Ph. Eur. / USP-NF / JP) and 25 % Cellulose Microcrystalline (Ph. Eur. / USP-NF / JP).

The compound lactose monohydrate produced at MEGGLE is not chemically synthesized. Organic solvents, catalysts and other reagents which might be a reason for the presence of secondary, tertiary, or quaternary amines and nitrite salts are not used. Furthermore, strongly acidic conditions as necessary prerequisite for the formation of nitrosamines are not achieved during manufacturing process. Lactose monohydrate is isolated and purified from whey which is a by-product of cheese manufacturing. Whey does not contain relevant amounts of nitrates (as precursor of nitrites) and nitrites. Testing on nitrates and nitrites is conducted as part of incoming goods inspection. The acceptance limits are as follows: Nitrate < 50 ppm, Nitrite < 5 ppm.

The compound cellulose microcrystalline is not chemically synthesized. It is of vegetable origin. Secondary and/or tertiary amines, amides, primary amines and ammonium salts are not used or present in the manufacturing process. The supplier certifies that the substance is free from nitrosamines.

The product MicroceLac® 100 is not chemically synthesized. Organic solvents, catalysts and other reagents which might be a reason for the presence of secondary, tertiary, or quaternary amines and nitrite salts are not used. Furthermore, strongly acidic conditions as necessary prerequisite for the formation of nitrosamines are not achieved during manufacturing process.

Test results on nitrites and nitrosamines are available.

Conclusion: Neither the chemical composition nor the processing conditions indicate any possibility for nitrosamine contamination and formation in the process or storage of the MEGGLE products.

Best regards

MEGGLE GmbH & Co. KG

Dr. Stefan Dreiheller

Regulatory Affairs / Specification Management

Nitrosamine Risk Evaluation

1) Applicable category based on structure and origin of the excipient in support to evaluate the risk of formation of nitrosamines in the excipient¹

Target Excipient:
Nitrogen containing?

➔ Yes

➔ No

Proteins, enzymes, products of fermentation or extraction of biologic sources, ...

Synthetic origin and nitrogen containing

Mined excipients, N-free products of fermentation or natural origin, ...

N-free mineral acids or bases, organic solvents, polymers, inorganic salts, small organic N-free entities, ...

➔ No

➔ Yes

Chemical Synthetic Manufacturing Process?
 including processes to introduce chemically synthesized fragments to biological products or substances of natural origin

2) Is sodium nitrite (NaNO ₂) or any other nitrite or nitrosating agent ² : <ul style="list-style-type: none"> - used in any steps in the manufacturing process³ as reagents/catalyst? - known to be used in the preparation of raw materials or intermediates used in the manufacturing process? - known to be used in the preparation of reagents/catalysts/processing aids used in the manufacturing process? - known or likely to be generated as impurities during the manufacturing process? - deliberately added to the process, including components of cell culture media or for fermentation? 	YES	NO	<i>Information not available</i>
	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<i>Information not available</i>
	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
3) Have you analysed the excipient for: <ul style="list-style-type: none"> - Nitrites? - Nitrosamines? Tests conducted on lactose monohydrate with other trade names in a contracted laboratory. / Nitrites: Method FIA; LoD 0.2 ppm / Nitrosamines NDMA, NDEA, NDPA, NDBA, NMOR, NPYR, NPIP: Method GC/TEA; LoQ 0.5 ppb	YES	NO	<i>Test result < LoD < LoQ</i>
	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<i>Test result < LoD < LoQ</i>
	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<i>Test result < LoD < LoQ</i>
4) Is water used in the manufacturing process? If "Yes": <ul style="list-style-type: none"> i. Is purified water⁴ used in the manufacturing process? ii. If potable water is used, where possible, please report the maximum level of nitrite. Maximum levels of <ul style="list-style-type: none"> - Nitrites 	YES	NO	<i>Not applicable</i>
	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<i>Not applicable</i>
	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
	_____		<i>Not available</i>
	<input type="checkbox"/>		<input type="checkbox"/>
5) Are there any secondary and/or tertiary amines ⁵ present in the manufacturing process as: <ul style="list-style-type: none"> - Raw material⁶? - Intermediate? - Reagent? - Processing aids? - Catalyst? - Solvent? If yes, are those amines present in the <ul style="list-style-type: none"> - Same - Previous - Subsequent step as any nitrosating agent mentioned in section 2?	YES	NO	<i>Not applicable</i>
	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<i>Not applicable</i>
	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<i>Not applicable</i>
	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<i>Not applicable</i>
	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<i>Not applicable</i>
	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<i>Not applicable</i>
	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<i>Not applicable</i>
	<input type="checkbox"/>	<input checked="" type="checkbox"/>	<i>Not applicable</i>
	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
	<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>

¹ Nitrogen-free materials are considered to be of lower inherent risk for nitrosamine contamination as they are typically manufactured and do not contain nitrosatable structures. Nitrosamines have been observed in medicinal products with N-containing APIs of chemical synthetic origin. EMA concludes that there is a very low risk of nitrosamines being present as impurities in biological medicinal products, although it can't be completely ruled out.

² See Guidance 1 in Annex

³ In this document, "manufacturing process" refers to the manufacturing steps that are outlined in the flow chart of the manufacturing procedure for the mentioned product.

⁴ Prepared by distillation, ion exchange, reverse osmosis

⁵ See Guidance 2 in Annex

⁶ 2020 IPEC General Glossary of Terms and Acronyms, <https://www.ipec-europe.org/glossary.html>



Nitrosamine Risk Evaluation

6) Is there any amide, primary amine or ammonium salt used or present in the substance manufacturing process as: - Raw material? - Intermediate? - Reagent? - Processing aids? - Catalyst? - Solvent? - Washing Fluid?	YES <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	NO <input checked="" type="checkbox"/> <input checked="" type="checkbox"/> <input checked="" type="checkbox"/> <input checked="" type="checkbox"/> <input checked="" type="checkbox"/> <input checked="" type="checkbox"/> <input checked="" type="checkbox"/>	
7) Recycled/recovered Solvents ⁷ : - Are recycled / recovered nitrogen containing solvents used in the manufacturing process?	YES <input type="checkbox"/>	NO <input checked="" type="checkbox"/>	
8) Equipment: - Is the substance produced in multipurpose equipment? - In case of multipurpose equipment, is the equipment used for manufacturing of any material involving nitrites, nitrosating agents or material with identified risk of formation of nitrosamines? - Are chloramines used as part of cleaning procedures used for manufacturing equipment?	YES <input checked="" type="checkbox"/> <input type="checkbox"/> <input type="checkbox"/>	NO <input type="checkbox"/> <input checked="" type="checkbox"/> <input checked="" type="checkbox"/>	Not applicable <input type="checkbox"/>
9) Additional comments, if any, not covered in the questionnaire			

Annex⁸:

Guidance 1 (Sources of nitrosating agents)

Nitrosating agents to be considered include: nitrites (e.g. sodium nitrite, NaNO₂) and nitrous acid (HNO₂), nitric oxide (NO), nitrosyl halides (e.g. ClNO, BrNO), dinitrogen trioxide (N₂O₃), dinitrogen tetroxide (N₂O₄) and organic nitrites (e.g. t-BuONO).

Other potential nitrosation risks:

- Side reaction in nitration reactions. Nitric acid typically contains nitric oxide as an impurity, additional nitrous acid may also be produced, leading to nitrosation, if any reducing agents are present.
- Hydroxylamine under oxidative conditions.
- Chloramines are known to generate N-nitrosamines under certain conditions and so should also be considered.⁹
- Ozone may lead to the formation of N-nitrosamines by initial oxidation of amines to nitrite.
- Use of azide salts and azide compounds is commonly followed by quenching with nitrous acid or nitrites and may lead to nitrite residues.
- Nitric acid and nitrates under reducing conditions may result in by-products with nitrosating activity.

This evaluation must include the use of all chemicals within a process, including those used during the quench and work-up as well as during reactive chemistry.

Guidance 2 (Sources of secondary and tertiary amines)¹⁰

Secondary amines are of greatest concern, however tertiary amines can also undergo nitrosation via more complex pathways. All secondary and tertiary aliphatic and aromatic amines should therefore be considered including those present as part of the starting material, intermediate or final structure as well as those introduced as reagents, catalysts, solvents or as impurities.

Tertiary amine bases (i.e. triethylamine, diisopropylethylamine and N-methylmorpholine) are known to degrade to secondary amines and have been implicated in N-nitrosamine formation.

Amines may also be introduced as impurities or degradants:

- Of common amide containing solvents such as N,N-dimethylformamide (DMF), N,N-dimethylacetamide (DMAC) and N-methylpyrrolidinone (NMP)
- Of quaternary ammonium salts such as tetrabutylammonium bromide (TBAB)
- Of primary amines such as monoethylamine
- Of starting materials, intermediates or the product itself

This evaluation must include the use of all chemicals within a process, including those used during the quench and work-up as well as during reactive chemistry.

Guidance 3 (Potential contamination risks)

Consider all potential sources of contamination in input materials.

Use of recovered materials (solvents, reagents, catalysts) is of particular concern if appropriate controls are not put in place. The materials DMF, ortho-xylene and tributyltin chloride were highlighted by the EMA as materials at risk of cross contamination by N-nitrosamines. Sodium azide was highlighted by Health Canada for risk of cross contamination with nitrite.

Cross contamination from other processes using shared equipment should be considered. Steps performed under GMP (using solvents/reagents with appropriate controls, and controls on their recovery and reuse) are considered to be a lower cross contamination risk.

⁷ See Guidance 3 in Annex

⁸ This information is partly transferred from the EFPIA decision tree for drug substances, published 1 Nov 2019

⁹ Nawrocki, J et al. Nitrosamines and Water, J. Hazard. Mater. 2011, 189, 1-18.

¹⁰ SCCS (Scientific Committee on Consumer Safety), Opinion on Nitrosamines and Secondary Amines in Cosmetic Products, 27 March 2012.